

Exponent 1/4 from the mean field approximation

Aleksey Nudelman

Department of Physics
University of California
Santa Barbara, CA 93106-9530 U.S.A.

email: anudel@physics.ucsb.edu

Abstract

We consider the magnetization of the three-dimensional ferromagnetic near the critical temperature in the mean field approximation. It is found that for some directions of easy magnetization one obtains exponent 1/4 instead of the usual 1/2.

KEY WORDS: Critical exponent. Spontaneous magnetization. Mean field approximation. Directions of easy magnetization.

Our model in three dimensions is as follows: we consider the cubic crystal lattice with spins situated at vertexes and introduce a coordinate system along the edges of the cell. Then directions of the spin can be specified relative to this basis as a unit vector (a_1, a_2, a_3) . To make our results coordinate-independent we induce the symmetry transformation around a vertex. We apply three 90-degree rotations C^4 around coordinate axes 1, 2, 3 and make one of the planes 12, 13, 23 a mirror plane. Doing this we generate from any given spin orientation 48 other directions (a_1, a_2, a_3) obtained by 3! permutations of a_1, a_2, a_3 times 8 permutations of the sign. (Compare it with the conventional mean field approach where spins are either up or down.) Magnetization per unit cell M in the direction of the external magnetic field H is defined as

$$M = \frac{\sum_{i=1}^{48} p_i \exp[\beta H p_i]}{Z[H]} \quad (1)$$

where p_i is a projection of the spin on the direction of the magnetization, β is the inverse temperature and i specifies a given spin orientation generated by the symmetry group. The statistical sum is defined as

$$Z[H] = \sum_{i=1}^{48} \exp[\beta H p_i] \quad (2)$$

In the mean field approximation the magnetization "eats its own tail" [1] i.e.

$$M = \frac{\sum_{i=1}^{48} p_i \exp[\beta M p_i]}{Z[M]} \quad (3)$$

This nonlinear equation can be solved near the critical temperature where the spontaneous magnetization becomes small. Given the fact that the right hand side of eq.3 is an odd function of M , the first three terms in the Taylor expansion are proportional to M , M^3 , M^5 respectively. If we are to limit ourselves to the first term of the Taylor expansion we would get an expression for the critical temperature in terms of p and k_B after M being factored out from both sides of eq.3. Then eq.3 would look like

$$1 \approx \frac{T}{T_c} + f(a_1, a_2, a_3) \frac{M^2}{T^2} + g(a_1, a_2, a_3) \frac{M^4}{T^4} \quad (4)$$

where

$$f(a_1, a_2, a_3) = p^2/6k_B^2 [(4a_2^2 a_3^2 - a_1^4)(a_2^4 + a_3^4) + (4a_1^2 a_3^2 - a_2^4)(a_1^4 + a_3^4) + (4a_1^2 a_2^2 - a_3^4)(a_1^4 + a_2^4)]$$

and $g(a_1, a_2, a_3)$ is to be specified later. Please note that we expressed the contributions of all 48 directions of the spins in terms of the first direction, (a_1, a_2, a_3) , from which they were generated by the symmetry group. Our focus now is on the different classes of the spins generated by (a_1, a_2, a_3) . For all practical purposes we can neglect the third term in eq.4 relative to the second term and the magnetization becomes

$$M \approx \frac{k_B T}{p} \sqrt{\frac{6(1 - T/T_c)}{f(a_1, a_2, a_3)}} \quad (5)$$

where $T_c = \frac{p^2}{3k_B}$. By the substitution

$$x = 1 - a_3^2 = a_1^2 + a_2^2 \quad (6)$$

$$a = \frac{a_1^2 a_2^2}{(a_1^2 + a_2^2)^2} \quad (7)$$

we obtain

$$M \approx \frac{k_B T}{p} \sqrt{\frac{3(1 - T/T_c)}{x[-5x^3(a-1)^2 + 10x^2(1-a) - x(7-2a) + 2]}} \quad (8)$$

This procedure would fail if $f(a_1, a_2, a_3)$ vanishes. If this happens we may no longer neglect the third term in eq.4 and one obtains $M \sim (T_c - T)^{1/4}$ instead of eq. 8. Let us obtain the range of a corresponding to the exponent 1/4. It follows from eqs. 6-7 that for $x \in [0, 1]$ and $a \in [0, 1/4]$ $f(a_1, a_2, a_3)$ vanishes iff $x = 0$ or $x = 1$. Please note that $x = 0$ corresponds to the orientation of the spins in the faces of the unit cell. Thus $x = 0$ is a particular case of $x = 1$. To illustrate this simple case we have plotted the dependence of the magnetization on the temperature in figure 1.




Figure 1: The dependence of the magnetization on the temperature when spins are aligned along the edges of the three dimensional cubic cell.

It is interesting to see if the exponent 1/4 obtained above is consistent with the mean field approximation. Let us introduce the two-particle correlation function $\langle \vec{p}_1 \vec{p}_2 \rangle / \langle \vec{p}^2 \rangle$ ($\langle p \rangle$ means statistical average of p and D is the dimension of the cubic lattice) for the simple case $x = 0$

$$\langle \vec{p}_1 \vec{p}_2 \rangle = \frac{\cosh[2Mp/k_B T] - 1}{1 + \cosh[2y] + 4(D-1)\cosh[Mp/k_B T] + 2(D-1)^2} \quad (9)$$

$$\langle \vec{p}^2 \rangle = 1 - \frac{D-1}{\cosh[Mp/k_B T] + D-1} \quad (10)$$

As it is seen from figure 2, the exponent 1/4 does not lead to any qualitative change in the behavior of the correlation function as compared to the well studied 1/2 case. Let us note that the mean field approximation is derived




Figure 2: The dependence of the two-particle correlation function on $Mp/k_B T$ for the 3D cubic crystal with the critical exponent $1/4$

under the assumption that the correlation between spins at the different lattice sites is small. As it is seen from the graph the correlation function remains small as long as the magnetization is close to the critical temperature. This behavior is to be contrasted with that of the 4D cubic crystal which we consider below.

To simplify calculations we consider the four-dimensional cubic lattice. A spin may align itself in 8 directions along the cell's edges. The magnetization

$$M = p \frac{\sinh[Mp/k_B T]}{\cosh[Mp/k_B T] + 3} \quad (11)$$

is no longer small near the critical temperature as can be seen from figure 3. The 4D case (as well as any higher dimensional lattice) is not described well by the mean field approximation because, as seen from figure 4, the two-particle correlation function is not small near the critical temperature.

In conclusion we show that the exponent $1/4$ is not possible in the two-dimensional case. (It is also not possible to get $1/4$ for the one-dimensional crystal as well but the calculations are so trivial that we do not show them here.) In the two-dimensional crystal we define the direction of easy magnetization as (a_1, a_2) . There are 8 directions of the spin relative to M generated




Figure 3: The dependence of the magnetization eq.11 on the dimensionless temperature.




Figure 4: The dependence of the two-particle correlation function on $Mp/k_B T$ for the 4D cubic crystal.

by the inversion around the coordinate origin and by the 90-degree rotations in the plane of the crystal. Near the critical temperature, after some algebra, one obtains the following expression for the magnetization:

$$M \approx \frac{2Tk_B}{p} \sqrt{\frac{1 - T/T_c}{1 - \frac{1}{3}(1 + (a_1^2 - a_2^2)^4 + (2a_1a_2)^4)}} \quad (12)$$

where $T_c = \frac{p^2}{2k_B}$ and p is a spin magnetic moment. If we substitute $a_1 = \cos\phi$ $a_2 = \sin\phi$ then $M \approx \frac{2Tk_B}{p} \sqrt{\frac{1 - T/T_c}{1 + \frac{\sin^2 4\phi}{2}}}$

$$1 + \frac{\sin^2 4\phi}{2} \neq 0 \quad (13)$$

and therefore

$$M \sim \sqrt{(T - T_c)} \quad (14)$$

for all directions of easy magnetization.

Acknowledgements

It is a pleasure to thank S. I. Ben-Abraham, Walter Poetz, Andreas X for discussions and Ms. Y. C. who worked hard on my grammar but refused to be acknowledged.

References

- [1] G.H.Wannier, Elements of Solid State theory, Cambridge University Press, 1959, p.90 .